

Calidad de vida relacionada con la salud, percepción de enfermedad, felicidad y emociones negativas en pacientes con diagnóstico de artritis reumatoide

Stefano Vinaccia Alpi^{*1}, Japcy Margarita Quiceno², Francy Lozano¹ y Sebastian Romero¹

¹ Fundación Universitaria SANITAS, Bogotá, Colombia. ² Universidad de Medellín, Antioquia, Colombia.

Recibido, febrero 22/2016

Concepto de evaluación, abril 17/2016

Aceptado, julio 26/2016

Referencia: Vinaccia, S., Quiceno, J.M., Lozano, F. & Romero, S. (2017). Calidad de vida relacionada con la salud, percepción de enfermedad, felicidad y emociones negativas en pacientes con diagnóstico de artritis reumatoide. *Acta Colombiana de Psicología*, 20(1), 60-69. DOI: 10.14718/ACP.2017.20.1.4

Resumen

El propósito de este estudio fue evaluar las relaciones entre la calidad de vida relacionada con la salud (CVRS), la percepción de enfermedad, la felicidad, la ansiedad y la depresión en 62 pacientes con diagnóstico de artritis reumatoide de la ciudad de Bogotá, Colombia. El diseño del estudio fue descriptivo correlacional; se utilizaron los instrumentos Escala de calidad de vida en artritis reumatoide (QOL-RA), Cuestionario de Percepción de Enfermedad (IPQ-B), Escala Subjetiva de Felicidad (SHS) y Escala Hospitalaria de Ansiedad y Depresión (HAD); y como resultados se encontró que hubo una favorable CVRS en estos pacientes posiblemente debido a la presencia de algunas dimensiones del QOL-RA, como el apoyo, la vida social y el estado de ánimo, los cuales han demostrado tener un impacto importante sobre la calidad de vida. Con el modelo de regresión lineal múltiple se encontró un peso negativo para la ansiedad y uno positivo para la felicidad con el QOL-RA. Se concluye que estos factores psicológicos negativos y positivos tienen un peso relevante sobre la percepción de calidad de vida de los pacientes con AR.

Palabras clave: Calidad de vida relacionada con la salud, percepción de enfermedad, felicidad, ansiedad, depresión, artritis reumatoide.

Health-related quality of life, illness perception, happiness and negative emotions in rheumatoid arthritis patients

Abstract

The aim of this study was to evaluate the relationship between quality of life related to health, illness perception, happiness, anxiety and depression in 62 patients diagnosed with rheumatoid arthritis. The study design was descriptive correlational. Instruments: The Quality of Life in Rheumatoid Arthritis Scale -QOL-RA, Brief Illness Perception Questionnaire -IPQ-B, Subjective Happiness Scale -SHS and The Hospital Anxiety and Depression Scale -HADS. Results: There was a high quality of life related to health in these patients, as some domains of QOL-RA such as support, social life and mood have demonstrated a significant impact on HRQOL. In the multiple linear regression model negative weight of anxiety and positive happiness with QOL-RA were appreciated. Conclusion: psychological factors such as anxiety and happiness have significant weight on the perceived quality of life of patients with RA.

Key words: Health-related quality of life, illness perception, happiness, anxiety, depression, rheumatoid arthritis.

* Fundación Universitaria Sanitas. Carrera 7 No. 173-64. Bogotá, Colombia. vinalpi47@hotmail.com

Qualidade de Vida Relacionada com a Saúde, Percepção de Doença, Felicidade e Emoções Negativas em Pacientes com Diagnóstico de Artrite Reumatoide

Resumo

O propósito deste estudo foi avaliar as relações entre a qualidade de vida relacionada com a saúde (QVRS), a percepção de doença, a felicidade, a ansiedade e a depressão em 62 pacientes com diagnóstico de artrite reumatoide da cidade de Bogotá, Colômbia. O estudo foi descritivo correlacional; utilizaram-se os instrumentos Escala de qualidade de vida em artrite reumatoide (QDV-RA), Questionário de Percepção de Doença (IPQ-B), Escala Subjetiva de Felicidade (SHS) e Escala Hospitalar de Ansiedade e Depressão (HAD); e como resultados verificou-se que houve uma favorável CVRS nestes pacientes possivelmente devido à presença de algumas dimensões do QOL-RA, como o apoio, a vida social e o estado de ânimo, os quais demonstraram ter um impacto importante sobre a qualidade de vida. Com o modelo de regressão linear múltipla verificou-se um peso negativo para a ansiedade e um positivo para a felicidade com o QDV-RA. Conclui-se que estes fatores psicológicos negativos e positivos têm um peso relevante sobre a percepção de qualidade de vida dos pacientes com AR.

Palavras-chave: Qualidade de vida relacionada com a saúde, percepção de doença, felicidade, ansiedade, depressão, artrite reumatoide.

INTRODUCTION

The definition of Rheumatoid Arthritis has been documented in the medical literature for more than two hundred years, but its etiology remains unknown (Entezami, Fox, Clapmam and Chung, 2011). Despite the fact that infectious agents such as viruses, bacteria and fungi, as well as environmental components like tobacco addiction, and even genetic factors have been regarded as causal factors, none of them has proven to be the cause of this disease (Julia and Marsal 2013; Pratt and Isaacs, 2015).

Rheumatoid Arthritis affects approximately 1% of the world's population, and the consequences for the person's health as the disease progresses are loss of joint mobility, deformity, disability, chronic pain and the presence of negative emotions such as anxiety, depression, stress and hopelessness (Gibofsky, 2012; Santiago, Geenen, Jacobs & Da Silva, 2015; Zhao, Chen & Chen, 2015), in addition to physical and emotional dependence, fatigue, morning stiffness and sleep disorders, among other ailments (Purabdollah et al., 2015).

In Colombia, epidemiological studies have estimated a prevalence of RA between 0.8 - 1.0% of the population. The average age of appearance of the disease is 53.2 years and for the onset of symptoms is 41.3 years, with higher prevalence in women (81.9%); co-morbidity is associated with essential hypertension, osteoporosis, and Sjogren's syndrome (Bautista et al., 2015; Machado, Ruiz & Medina, 2015).

Rheumatoid Arthritis therefore implies a major change in the health-related quality of life (HRQOL) for people who suffer from it (Matcham, Scott, Rayner & Hotopf; Akiskal,

2014). HRQOL is the assessment that a person makes, in accordance with its own criteria, of their physical, emotional and social state at a given time, and reflects the degree of satisfaction with a personal situation at a physiological level (general symptomatology, functional disability, analytical situation, sleep, sexual response) emotional level (feelings of sadness, fear, insecurity, frustration) and social level (work or school situation, social interactions social in general, family relationships, friendships, economic level, participation in the community, leisure activities, among others) (Schwartzmann, 2003).

The person with a diagnosis of Rheumatoid Arthritis must face, not only the symptoms associated with the disease, but the ambiguity of these (Purabdollah et al., 2015). Since Rheumatoid Arthritis is one of the main conditions that most diminishes quality of life (Garip, Eser & Bodur, 2011). However, the prevalence of symptoms in situations of physical and functional deterioration does not exclude the presence of positive emotions such as happiness, which serve as protective frames for the physical and mental health of patients with chronic medical conditions (Angner, Ray, Saag & Allinson, 2009; Angner, Gandhi, Purvis, Amante and Allinson, 2013; 2015).

According to Palomera (2009) "happiness is composed of three dimensions: positive emotions, negative emotions and vital satisfaction." "Happiness is the result of the degree in which the experiences of positive affectivity exceed experiences of negative affectivity and the degree in which a person perceives that his/her goals and motivations are resolved (p. 273)". In addition, the ability to maintain positive emotionality during periods of illness has also been associated with a better welfare (Lyubomirsky, Sheldon &

Schkade, 2005; Steptoe & Wardle, 2005). The literature is consistent in affirming the positive relationship between quality of life and happiness in health (Musschenga, 1997)

On the other hand, cognitive and emotional factors play an important role in the perception the patient has of his disease, influencing the development and course of the same. The perception of the disease refers to a cognitive and emotional representation that the patient makes of his condition, whereby if this implies a threat to his/her health, the representation orients the actions and the use of coping strategies to deal with the disease (Quiceno & Vinaccia, 2010).

In this regard, the perception of the disease is related to a number of outcome measures in Rheumatoid Arthritis, including disability, low mood, pain and decline in physical function (Prajapati et al., 2014). It can even delay the search for medical treatment, which may exacerbate the symptoms of the disease and lead to long-term loss of several functions of daily life such as the ability to travel, personal care, work capacity, financial independence and participation in social roles, among other basic activities (Van der Elst et al., 2015). In different studies, a directly proportional relationship between HRQOL and the perception of illness has been found (Quiceno & Vinaccia, 2010).

According to the above, the aim of this study was to evaluate the relationships between quality of life related to health, the perception of disease, happiness, anxiety and depression in 62 patients with a diagnosis of rheumatoid arthritis, residents in the city of Bogotá.

METHOD

Participants

62 adult patients (men and women) participated in this study with a diagnosis of rheumatoid arthritis according to the classification criteria for this disease of the year 2010. The sample was selected through a non-random sampling of the available subjects, who attended a foundation of the city of Bogotá, which provides support to the rheumatic patients. Inclusion criteria were being 18 years or older, not having any diagnosis of cognitive impairment and accepting freely and voluntarily to participate in the the investigation.

Instruments

Quality of Life in Rheumatoid Arthritis Scale, QOL-RA
The QOL-RA is a specific questionnaire that evaluates the HRQOL of patients with arthritis Rheumatoid. It was originally developed and validated to Spanish by Danao, Padilla and Johnson (2001). Cronbach alpha coefficients for this

instrument range from 0.87 to 0.90. The Colombian version by Vinaccia, Riveros, Quiceno, and Anaya was used in this research (in press). This scale contains eight items where each one constitutes a dimension, thus a Cronbach Alpha analysis is not carried out. Its eight dimensions are: physical ability, support, pain, stress, health, arthritis, social life and mood; in addition it contains a full scale. The response system is Likert type ranging from 0 (very poor quality of life) to 10 points (excellent quality of life), where the higher the score, the better perception of HRQOL. Cronbach's alpha coefficient for the total scale for this study was 0.949.

The Brief Illness Perception Questionnaire, IPQ-B

Originally developed by Broadbent, Petrie, Main and Weinman (2006). It contains eight items where each one makes up a dimension, and therefore a Cronbach Alpha analysis is not conducted. The response system is Likert type ranging from 0 to 10 points. The dimensions are: consequences, duration, personal control, treatment control, identity, worry, emotional response and understanding of the disease.

Subjective Happiness Scale, SHS

Developed originally by Lyubomirsky and Lepper (1999), possesses a high level of reliability ($\alpha < .80$). It is comprised by four items that are answered through a Likert system that goes from 1 to 7 points, where the higher the score, the better the subjective perception of happiness. The Cronbach's alpha coefficient for this study was .711.

The Hospital Anxiety and Depression Scale, HADS

The Colombian version of the HADS scale was developed by Rico, Restrepo and Molina (2005) with a Cronbach's alpha coefficient of .85. It includes 14 items that are answered in a Likert type system with 4 response options ranging from 0 to 3 points, and has two dimensions that evaluate anxiety and depression related to the current period. The range for each dimension goes from 0 to 21 points, where the higher the score, the higher the level of anxiety or depression. Cronbach's alpha coefficient for this study was 0.824 for anxiety and 0.795 for depression.

Procedure

After obtaining the consent and acceptance from the managers and the Scientific Committee of the foundation that supported the research, patients were invited to participate in the study. Those who accepted the invitation were assembled on the premises of the foundation in groups of maximum 20 people. The purposes and scope of the research were clarified to each group in order for them to sign the informed consent freely and voluntarily. Later, in a self

administrated manner and counting with the help of the researchers in charge of the study when patients required it, each patient answered the questionnaires (QOL-RA, IPQ-B, SHS and HADS) and a socio-demographic data sheet. Application time for each participant was one hour on average.

RESULTS

The results of the study were analyzed through the statistical package SPSS version 22. For the descriptive analysis of the variables age, time of diagnosis, happiness, anxiety and depression, perception of illness and HRQOL, the average was used as a measure of central tendency. Likewise, the standard deviation (D.T. =1.17) and the minimum (min.) and maximum (max.) scores of the different dimensions of the psychometric instruments used were calculated. With regard to nominal variables (sex, marital status, educational level, socio-economic status, occupation, with whom he/she lives, health system, taking specific medications for the disease or for psychiatric ones) percentages were used.

Additionally, the Cronbach's alpha coefficient was used for the reliability analysis of the psychometric instruments applied in the study. The Pearson's r coefficient was calculated for the correlation analysis between age and time of diagnosis (demographic variables) and for the different dimensions of the questionnaires of quality of life related to health (QOL-RA, IPQ-B, SHS and HADS), having previously calculated the distribution of variables (Kolmogorov-Smirnov). Finally, a multiple linear regression model was employed considering the selection method by steps to analyze the weight of the predictor variables such as age, time of diagnosis, happiness, anxiety and depression and perception of disease on HRQOL (criterion variable)

Descriptive Analysis according to sociodemographic characteristics

Table 1 presents the descriptive data of the socio-demographic and clinical variables of the research. In this study, a prevalence of the female sex over the male can be appreciated, with an average age of 59.7* years and a range between 32 to 79 years. There was a predominance of the sample in the categories *married* and *separated/divorced* according to marital status. Most of them achieved an educational level of basic secondary (high school) and belong to a Colombian medium low socioeconomic status (level 1, 2 and 3). Similarly, a great portion of the sample were retired from their working life and a smaller percentage was dedicated to household activities. In addition, a large proportion of the participants lived with relatives and there were very few who lived on

their own. With respect to clinical variables, it is observed that patients had 17 years average of having been diagnosed with AR, within a range from 2 to 60 years. Most were covered by a health insurance system (health promotion company), did not take psychiatric medication but did follow a specific treatment for their disease.

Table 2 presents the descriptive data obtained from the instruments used in table 2.

At a general level, there was a level of reliability, from acceptable to very good, in different dimensions and full scales of the psychometric instruments used in the study as shown in table 2.

Regarding the descriptive results shown in table 2, it is observed that *happiness* (SHS), had an average score. With respect to the HADS scale, according to the criteria of Rico et al. (2005), in order to indicate if patients in Colombia present these symptoms or not, this study shows that the *anxiety dimension* had a score that limited with the cut-off point of 8 for the *anxiety* sub-scale, while *depression* had a low score with respect to the cut-off point of 9 for the *depression* sub-scale. In relation to the perception of disease (IPQ-B), the highest average scores were in the dimensions of *duration*, *control of treatment* and *understanding of the disease*, while the dimensions of *consequences*, *personal control*, *identity*, *worry* and *emotional response* had a moderate score. As for the QOL-RA of the HRQOL scale, the highest average scores were obtained in the dimensions of *support*, *social life and mood*, and average scores in *physical ability*, *pain*, *stress*, *health*, *arthritis*, *social life*, *mood* and in the QOL-RA full scale.

Pearson correlation analysis

Table 3 presents the results of the Pearson correlation analysis *between age, time of diagnosis, (MSM) happiness, anxiety depression (HAD), perception of disease (IPQ-B) with a HRQOL (QOL-RA)*, where a statistically significant relationship between some variables of the study was found at the 0.01 and 0.05 levels.

With regard to the Happiness Scale (SHS), moderate and high positive correlations with all dimensions of the questionnaire of CVRS QOL-RA were found, such as *physical ability*, *support*, *pain*, *stress*, *health*, *arthritis*, *social life and mood*, as well as with the full scale. Similarly, the anxiety and depression scale HAD had moderate to high negative correlations with all dimensions of HRQOL.

With respect to the Perception of Disease Questionnaire (IPQ-B), this instrument shows that the dimensions of *personal control*, *identity and concern* had low and moderate negative correlations with all dimensions of HRQOL. On the other hand, the dimension of *consequences* had low

Table 1
Sociodemographic characteristics of the sample (n=62)

Sociodemographic Factor					
	n	%		n	%
<i>Sex</i>			<i>Age</i>		
Man	3	4.8	Mean (D.T)	59.7	9.3
Woman	59	95.2	Mín.-Máx.	32	79
<i>Marital status</i>			<i>Educational level</i>		
Single	12	19.4	Primary	15	24.2
Married	19	30.6	High School	26	41.9
Divorced	19	30.6	Tecnology	11	17.7
Free union	5	8.1	Academic	7	11.3
Widower	7	11.3	Posgraduate	3	4.8
<i>Socioeconomic stratum</i>			<i>Occupation</i>		
1	2	3.2	Employed	4	6.5
2	16	25.8	Independ worker	7	11.3
3	29	46.8	Retired	30	48.4
4	13	21.0	Unemployed	2	3.2
5	1	1.6	Home	19	30.6
6	1	1.6			
<i>Who does he/she live with</i>					
Alone	6	9.68			
With family	56	90.32			
Clínical factors					
	n	%		n	%
<i>Health System</i>			<i>Diagnosis time</i>		
SISBEN	5	8.1	Mean (D.T)	205.7	170.3
EPS	48	77.4	Mín.-Máx.	23	720
Prepaid medicine	8	12.9			
Prívate	1	1.6			
<i>Taking psychiatric drugs</i>			<i>Get specific treatment for the disease</i>		
Yes	13	21.0	Yes	53	85.5
Not	49	79.0	Not	9	14.5

Nota: n= Number of participants; D.T= Standard deviation; Mín.= Mín.; Máx.= Máx.

Table 2
Descriptive statistics of happiness, anxiety, depression, perception of illness and HRQOL variables

	Mean	DT	Mín.	Máx.
<i>SHS: Happiness</i>	4.06	1.02	2	6
<i>HAD: Anxiety</i>	8.34	4.11	0	18
<i>HAD: Depression</i>	5.69	3.60	0	14
<i>IPQ-B: Perception of illness</i>				
Consequences	6.08	2.52	0	10
Timeline	8.31	2.49	0	10
Personal control	6.63	2.63	0	10
Treatment control	7.65	2.33	0	10
Identity	6.06	2.37	0	10
Concerned	6.34	2.69	0	10
Coherence	7.98	1.87	0	10
Emotional representation	6.45	2.86	0	10
<i>QOL-RA: CVRS in arthritis</i>				
Physical ability	6.40	2.38	1	10
Support	7.39	2.25	1	10
Pain	5.89	2.61	1	10
Tensión	6.77	2.66	1	10
Health	6.16	2.34	1	10
Arthritis	6.06	2.60	1	10
Interaction	7.13	2.53	1	10
Mood	7.05	2.43	1	10
QOL-RA Scale	6.62	2.13	1	10

Note: *n*= number of participants; *D.T*= Standard deviation; *Mín.* = Mín.; *Máx.*= Máx.

and moderate negative correlations with all dimensions of HRQOL except with *nervous tension*. Similarly, the *emotional response* dimension had low and moderate negative correlations with all dimensions of HRQOL except with the dimension of *pain*. Finally, the control of treatment dimension, of the IPQ-B had a low negative correlation with the *mood* dimension of HRQOL.

Multiple linear regression analysis

According to results found in the final models of Multiple Linear Regression, and considering the CVRS as dependent variable (DV), it was found that the CVRS (QOL-RA) presented a determination coefficient $R^2c =$

49.4 %, explained negatively by the *anxiety* dimension of the HAD Scale ($\beta = -0.453$; $t = -4.241$; $p = .000$), and positively for the Happiness Scale, SHS ($\beta = 0.364$; $t = 3.411$; $p = .001$).

DISCUSSION

In this study, the majority of participants were women with an average age of nearly 60 years, a secondary education level, a marital status either married or divorced / separated, who had retired from work, belonging to a medium / low Colombian socioeconomic stratum and who, in average, had a diagnosis of Rheumatoid Arthritis for 16 years and 6 months.

It is worth mentioning that some of the instruments used in this study have not been validated in the Colombian context. There are only characterizations of the instruments IPQ-B Perception of Illness (Vinaccia, Quiceno and Remor, 2012) and the Subjective Happiness Scale - SHS (Quiceno et al., 2012, 2013), whereas on the other hand, there is a Colombian validation of the Hospital Anxiety and Depression Scale HADS (Rico et al, 2005) and of the Quality of Life in Rheumatoid Arthritis Scale QOL - RA (Vinaccia, Riveros, Quiceno and Anaya, in evaluation).

Regarding the descriptive results of the study in accordance with the *perception of disease* (IPQ-B), the patients in this study know that their illness has a long-term prognosis and understand what it means; report symptoms and signs of their condition which limit them in some way in their daily lives; perceive that they have a moderate control of the medical treatment, and refer that they try to manage their disease to the best of their abilities. On the other hand, they perceive that the disease somehow affects their lives, experience some concern about the consequences the disease may cause them at the physical, social, economic and emotional level, and express emotional responses such as anger, annoyance, fear and sadness.

With regard to *happiness* (SHS), the participants in this study had lower scores ($M = 4.06$) compared to other researchers developed in Colombia using the same scale with samples of university students ($M = 5.0$) (Quiceno et al., 2013) and with men deprived of freedom ($M = 5.0$) (Quiceno et al., 2012). In this regard, in the patients with RA of this study, happiness levels are not so favorable compared to Colombian healthy samples, although different studies affirm that the Colombian population has very high rates of happiness and optimism, regardless of the negative or positive events that happen to them (CENEC, 2001; Cross & Torres, 2006; Sinisterra, 2007).

Table 3

Pearson correlation analysis between age, diagnosis time, happiness, anxiety, depression, perception of illness and HRQOL

	QOL-RA								QOL-RA Scale
	Physical ability	Support	Pain	Tension	Health	Arthritis	Support	Mood	
Age	-.054	.112	.061	-.067	-.018	.044	.155	-.116	.016
Diagnosis time	.009	.185	.1	.235	-.015	.094	.044	.078	.106
SHS: Happiness	.433**	.515**	.420**	.563**	.505**	.439**	.623**	.640**	.601**
HAD Anxiety	-.492**	-.454**	-.489**	-.581**	-.624**	-.510**	-.593**	-.674**	-.644**
HAD Depresión	-.489**	-.493**	-.410**	-.549**	-.509**	-.475**	-.481**	-.666**	-.591**
IPQ-B: Consequences	-.467**	-.442**	-.262*	-.234	-.432**	-.298*	-.440**	-.369**	-.424**
IPQ-B: Timeline	-.071	.04	.116	.124	-.037	.098	.001	.016	.044
IPQ-B: Personal control	.437**	.355**	.422**	.314*	.431**	.325**	.353**	.545**	.458**
IPQ-B: Treatment control	.171	.102	.184	.106	.227	.16	.191	.295*	.209
IPQ-B: Identity	-.530**	-.431**	-.366**	-.267*	-.477**	-.398**	-.419**	-.364**	-.470**
IPQ-B: Concemed	-.312*	-.334**	-.312*	-.395**	-.403**	-.345**	-.375**	-.417**	-.424**
IPQ-B:	-.002	.107	.11	.003	.072	-.01	.111	.141	.073
IPQ-B: Emocional representation	-.474**	-.403**	-.215	-.353**	-.421**	-.328**	-.480**	-.468**	-.455**

*p < .05, two tails

**p < .01, two tails

On the other hand, considering the levels of anxiety and depression (HADS), at a general level there were no symptoms of clinical depression in these patients (Rico et al., 2005), while there were in anxiety, since scores were slightly above the cut-off point for the Colombian sample (M = 8.34). These results differ in relation to the symptoms of depression in patients from the study of Matcham et al, 2014, but are similar to the study of Ryan (2014) in levels of anxiety in patients with RA.

In accordance with the HRQOL (QOL-RA), the patients of this study had higher average scores in the dimensions of *support, social life and mood*, and average scores in the dimensions of *physical ability, pain, nervous tension,*

health, arthritis, social life and, mood, and in the full scale of QOL-RA. The results of the full scale of the QOL-RA in this study with patients natives of the city of Bogotá (M = 6.62) are slightly similar to those obtained in studies of Vinaccia, Fernandez, Moreno and Padilla (2006) (M = 6.40), and differ a little from the study of Vinaccia, Tobon, Moreno, Cadena and Anaya (2005) (M = 5.28) with samples of AR natives of the “paisa culture” of Medellín. It can be seen that, despite the differences in the stage of the disease and culture in these three studies, the levels of HRQOL are kept almost similar. In other words, the cultural differences in this type of disease are not so marked in Colombian samples, which differs from other studies that have suggested

Table 4

Multiple linear regression analysis

	Unstandardized		Standardized coefficients		
	B	Standard error	Beta	t	Sig.
(Constant)	5.507	1.224		4.498	.000
Anxiety -HAD	-.235	.055	-.453	-4.241	.000
Happiness -SHS	.758	.222	.364	3.411	.001

the influence of ethnic and cultural factors in the perception of the quality of life both in the general population as in samples with any chronic disease (Goh, Rusli and Khalid, 2014; Verhagen, Ros, Steunenbergh and Witt, 2014).

In terms of correlation analysis, it can be seen that for these patients there were no significant correlations between *age* and *time of diagnosis of the disease* with an HRQOL. In terms of the Perception of Disease Scale (IPQ-B), the dimensions of *life, control of treatment and understanding of disease* had no correlations with the HRQOL. However, the other dimensions as *consequences, identity, worry and emotional response* were negatively associated with different dimensions and the full scale of HRQOL (QOL-RA), while the *personal control* dimension had a positive correlation with the HRQOL. In this regard, Quiceno and Vinaccia (2010) found in a review of literature with different physical pathologies that the *perception of disease* correlated with quality of life, demonstrating that the negative perceptions of the disease have a direct impact on the physical and mental well-being of people with any chronic condition (Fowler and Baas, 2006;) Groarke, Curtis, Coughlan and Gsel, 2004; Paschalides et al., 2004).

Similarly, it has been found that the HADS scale, that measures anxiety and depression, had negative correlations with all dimensions and with the full scale of QOL-RA. Different studies have shown revealed significant levels (21% to 70%) of anxiety in RA (Covic et al., 2012; Isik, Koca, Ozturk and Mermi, 2007) and the influence of this negative emotion on HRQOL (Nas et al., 2011; Ozcetin et al., 2007).

Positive correlations with all dimensions and the full scale of QOL-RA can be seen in relation to the Happiness Scale (SHS). In this regard, the research on happiness as subjective well-being construct has been investigated in health psychology as an important indicator of quality of life in samples of chronic patients in hemodialysis process (Bennet, Weinberg, Bridgman and Cummins, 2015; Musschenga, 1997), in elderly people with different medical diagnoses (Steptoe, Deaton and Stone, 2015) and schizophrenia (Palmer, Martin, Deep, Glorioso and Cesky, 2014).

On the other hand, multiple linear regression analysis shows that the anxiety dimension of the HADS scale had a negative weight on the quality of life, while the happiness of the SHS had a positive weight on quality of life. Similar results have been found in samples of chronically ill individuals (Hoppe, 2013 ;) NAS et al., 2011; Ozcetin et al., 2007; Veenhoven, 2008) and in patients with chronic pain (Müller et al., 2015; Tran et al., 2015; Ryan and McGuire, 2015).

Finally, as limitations of the study, there is, first of all, the limited size of the sample (62 people), which was

conditioned to the pathological characteristics (diagnosis of rheumatoid arthritis) of the study participants; secondly, the absence of solid epidemiological studies about this disease in Colombia and about the population with rheumatoid arthritis of the city of Bogota; and thirdly, the lack of instruments validated for the population under study.

The results of this research demonstrate the importance of both the salutogenic variables (happiness) and the pathogenic ones (anxiety, depression) on HRQOL of patients with rheumatoid arthritis. The diagnosis of a chronic illness such as RA profoundly affects the lives of people who suffer from them, since their life project has to be reviewed and often they may not know what its course will be, given that the characteristics of evolution are often unpredictable (Vinaccia, Ramirez & Toro, 2001). Several investigations have found in these patients different types of deficits in learned hopelessness (both at a motivational level, for the reduction in daily living activities; and at a cognitive level for the reduction of efforts to develop new adaptation strategies), that affect the emotional area with increased depression and anxiety, and diminished self-esteem (Vinaccia, Cadena, Juárez, Contreras & Anaya, 2004).

Considering that there is vast empirical information on the protector role of positive psychosocial variables and positive emotions for a better psychological and physical well-being in patients with RA (Quiceno and Vinaccia, 2011), it is necessary to develop programs for the promotion of positive emotions, resilience and training of self-control techniques in RA patients (Quiceno, Vinaccia and Remor, 2011; Gräninger, 2015).

REFERENCES

- Angner, E., Ray, M., Saag, K., & Allison, J. (2009). Health and happiness among older adults. *Journal of Health Psychology, 14*(4), 503-512.
- Angner, E., Ghandi, J., Purvis, K., Amante, D., & Allison, J. (2013). Daily functioning, health status and happiness in older adults. *Journal of Happiness Study, 14*, 1563-1574.
- Bautista, W., Fernandez, D., Jiménez, R., Cardozo, R., Marin, A., Soler, M. P., et al. (2015). Perfil epidemiológico de pacientes colombianos de artritis reumatoide evaluados en una clínica de especializada en atención integral. *Reumatología Clínica*. Doi.org/10.1016/j.reuma.2015.11.009
- Bennett, P. N., Weinberg, M. K., Bridgman, T., & Cummins, R. A. (2015). The Happiness and Subjective Well-Being of people on Haemodialysis. *Journal Renal Care, 5*(1). Doi: 10.1111/jorc.12116
- Broadbent, E., Petrie, K. J., Main, J., & Weinman, J. (2006). The Brief Illness Perception Questionnaire. *Journal of Psychosomatic Research, 60*, 631-637.

- Cruz, J. & Torres, J. (2006). "¿De qué depende la satisfacción subjetiva de los colombianos?" *Cuadernos de Economía*, 45, 131-154.
- CENEC (2001). *El enigma de ser colombiano*. Recuperado de: [http://: Semana.com](http://Semana.com): http://www.semana.com/wf_InfoArticulo.aspx?IdArt=93494
- Covic, T., Cummings, S. R., Pallant, J. F., Manolios, N., Emery, P., Conogham, P. G., et al. (2012). Depression and anxiety in patient's whit rheumatoid arthritis: prevalence rates on a comparison of depression, anxiety and stress scale (DASS) and the hospital, anxiety and depression scale (HADS). *BMC Psychiatry*, 24, 12-16.
- Danao, L. L., Padilla, G. V., & Johnson, D.A. (2001). An English and Spanish quality of life measure for rheumatoid arthritis. *Arthritis & Rheumatology*, 45, 167-173.
- Entezami, P., Fox, D., Clapmam P., & Chung, K. (2011). Historical Perspective on the Etiology of Rheumatoid Arthritis. *Hand Clinics*, 27, 1-10.
- Fowler, C. & Baas, L. S. (2006). Illness Representations in Patients with Chronic Kidney Disease on Maintenance Hemodialysis. *Nephrology Nursing Journal*, 33, 173-187.
- Garaigordobil, M. (2015). Predictor variables of happiness and its connection with risk and protective factors for health. *Frontiers in Psychology*, 6, 1176. Doi: 10.3389/fpsyg.2015.01176
- Graninger, M. (2015). Behavioral training as additional therapy Approach for rheumatoid arthritis. *Zeitschrift für Rheumatologie*, 74(7), 579-583.
- Garip, Y., Eser, F., & Bodur, H. (2011). Health-related quality of life in rheumatoid arthritis: comparison of RAQoL with other scales in terms of disease activity, severity of pain, and functional status. *Rheumatology International*, 31(6), 769-772.
- Gibofsky, A. (2012). Overview of epidemiology, pathophysiology, and diagnosis of rheumatoid arthritis. *American Journal Management Care*, 18(13), 295-302.
- Goh, S. G., Rusli, B. N., & Khalid, B. A. (2014). Diabetes quality of life perception in a multiethnic population. *Quality Of Life Research*, 24(7), 1677-1686.
- Groarke, A., Curtis, R., Coughlan, R., & Gsel, A. (2004). The role of perceived and actual disease status in adjustment to rheumatoid arthritis. *Rheumatology*, 43(9), 1142-1149.
- Julia, A. & Marsal, S. (2013). The genetic architecture of rheumatoid arthritis: from susceptibility to clinical subphenotype associations. *Current Topics in Medicine Chemistry*, 13(6), 720-731.
- Hoppe, S. (2013). Chronic Illness as a source of happiness. *Health, Culture and Society*, 5(1). Doi 10.5195/hcs.2013.138
- Isik, A., Koca, S. S., Ozturk, A., & Mermi, O. (2007). Anxiety and depression in patients with rheumatoid arthritis. *Clinical Rheumatology*, 26(6), 872-878.
- Lyubomirsky, S. & Lepper, H. (1999). A measure of subjective happiness: Preliminary reliability and construct validation. *Social Indicators Research*, 46, 137-155.
- Lyubomirsky, S., Sheldon, K. M., & Schkade, D. (2005). Pursuing happiness: the architecture of sustainable change. *General Review Psychology*, 9(2), 111-131.
- Machado, J. E., Ruiz, A. F., & Medina, D. A. (2015). Epidemiología de la artritis reumatoide en una cohorte de pacientes colombianos. *Revista Colombiana de Reumatología*, 22(3), 148-152.
- Matcham, F., Scott, I., Rayner, L., & Hotopf, M. (2014). The Impact of rheumatoid arthritis the SF36. A Systematic review Meta-analysis. *Rheumatology*, 52(12), 2136-2148.
- Muller, R., Terrill, A. R., Jensen, M. P., Molton, I. R., Raveslot, C., & Ipsen, C. (2015). Happiness, pain intensity, pain interference, and distress in individuals with physical disabilities. *American Journal of Physical Medicine & Rehabilitation*, 94(12), 1041-1051.
- Musschenga, A. W. (1997). The relation between concepts of Quality-of-Life, Health and Happiness. *The Journal of Medicine and Philosophy*, 22(1), 11-28. Doi: 10.1093/jmp/22.1.11
- Nas, K., Sarak, A. J., Gur, A., Cevik, R., Altay, Z., Erdal, A., et al. (2011). Psychological status in associated Whit Health related quality of life in patients with rheumatoid arthritis. *Journal Back Musculoskelet Rehabilitation*, 24(2), 95-100.
- Oz Cetin, A., Ataoglu, S., Kocer, E., Yazici, S., Yaldiz, O., Ataoglu, A., et al. (2007). Effects of depression and anxiety on quality of life of patients with rheumatoid arthritis, knee osteoarthritis, fibromyalgia syndrome. *Western Indian Medicine Journal*, 56(2), 122-129.
- Palmer, B. W., Martin, A. S., Deep, C. A., Glorioso, D. K., & Jeste, V. (2014). Wellness whit Wilness: Happiness in schizophrenia. *Schizophrenia Research*, 159(1), 151-156.
- Palomera, R. (2009). Educando para la felicidad. En E.G. Fernández-Abascal (Ed.), *Emociones positivas* (pp. 247-274). Madrid, España: Pirámide.
- Pascalides, C., Wearden, A. J., Dunkerley, R., Bundy, C., Davies, R., & Dickens, C. M. (2004). The associations of anxiety, depression and personal illness representations with glycaemic control and health-related quality of life in patients with type 2 diabetes mellitus. *Journal Psychosomatic Research*, 57(6), 557-564.
- Prajapati, R., Plant, D., Maskell, D., Morgan, C., Ali, F., Morgan, A., Wilson, A., et al. (2014). Impact of psychological factors on subjective disease activity assessments in patient severe whit severe rheumatoid arthritis. *Arthritis Care Research*, 66(6), 661-668.
- Pratt, A. G. & Isaacs, J. D. (2015). Genotyping in rheumatoid arthritis: a game changer in clinical management? *Expert Review Clinical Immunology*, 11(3), 303-305.
- Purabdollah, M., Lakdizaji, S., Rahmani, A., Hajaliu, M., & Anzarin, K. (2015). Relationship between sleep disorders,

- pain and quality of life in patients with rheumatoid arthritis. *Journal of Caring Sciences*, 4(3), 233-241.
- Quiceno, J. M. & Vinaccia, S. (2010). Percepción de enfermedad: Una aproximación a partir del Illness Perception Questionnaire. *Revista Psicología desde el Caribe*, 25, 56-83.
- Quiceno, J. M. & Vinaccia, S. (2011). Evolución de la intervención psicológica en artritis reumatoide. *Revista Psicología desde el Caribe*, 27(1), 160-178.
- Quiceno, J. M., Vinaccia, S., Barrera, R., Latorre, R., Molina, D. C., & Zubieta, F. A. (2013). Resiliencia, felicidad, depresión, ideación suicida y afrontamiento del estrés en estudiantes universitarios colombianos. En J. Gaxiola & J. Palomar (Coords.), *Estudios de Resiliencia en América Latina (Vols. 2)* (pp. 71-88). México, D.F.: Pearson
- Quiceno, J. M., Vinaccia, S., Ibáñez, J., Álvarez, A., Jiménez, J., Pinzón, L., & Serna, A. M. (2012). Calidad de vida relacionada con la salud, resiliencia y felicidad en hombres privados de la libertad. *Pensamiento Psicológico*, 10(2), 23-33.
- Quiceno, J. M., Vinaccia, S., & Remor, E. (2011). Programa de potenciación de la resiliencia para pacientes con artritis reumatoide. *Revista de Psicopatología y Psicología Clínica*, 16(1), 27-47.
- Rico, J. L., Restrepo, M., & Molina, M. (2005). Adaptación y validación de la escala hospitalaria de ansiedad y depresión (HAD) en una muestra de pacientes con cáncer del Instituto Nacional de Cancerología de Colombia. *Avances en Medicina*, 3(1), 73-86.
- Ryan, S. (2014). Psychological effects of living with rheumatoid arthritis. *Nursery Standard*, 29(13), 52-59.
- Ryan, S. & McGuire, B. (2015). Psychological predictors of pain severity, pain interference, depression and anxiety in rheumatoid arthritis patients with chronic pain. *British Journal of Health Psychology*. Doi: 10.1111/bjhp.12171
- Santiago, T., Geenen, R., Jacobs, J. W., & Da Silva, J. A. (2015). Psychological factors associated with response to treatment in rheumatoid arthritis. *Current Pharmaceutical Design*, 21(2), 257-269.
- Schwartzmann, L. (2003). Calidad de vida relacionada con la salud: aspectos conceptuales. *Ciencia Enfermería*, 2, 9-21.
- Sinisterra, D. (2007). *Colombian happiness: A look at life satisfaction and explanatory style. Disertación Doctoral no publicada*. Florida Atlantic University, Boca Raton, FL, EE.UU.
- Stephoe, A. & Wardle, J. (2005). Positive affect and biological function in everyday life. *Neurobiology Aging*, 26(1), 108-112.
- Stephoe, A., Deaton, A., & Stone, A. (2015). Subjective Well-being, Health, and Ageing. *Lancet*, 385, 640-648.
- Tran, P., Nilakantan, A., Foote, D., Sturgeon, D., Mackey, S., & Johnson, K. (2015). Pain interference mediates the relationship between pain and happiness. *Journal of pain*, 16(4), 20. Doi: <http://dx.doi.org/10.1016/j.jpain.2015.01.091>
- Van der Elst, K., De Cock, D., Vecoven, E., Arat, S., Meyfroidt, S., Joly, J., Moons, P., et al. (2015). Are illness perception and coping style associated with the delay between symptom onset and the first general practitioner consultation in early rheumatoid arthritis management? An exploratory study within the Care RA trial. *Scandinavian Journal of Rheumatology*, 23, 1-8.
- Veenhoven, R. (2008). Healthy happiness. *Journal of Happiness Studies*, 9, 449-469.
- Verhagen, I., Ros, W., Steunenbergh, B., & DeWitt, N. (2014). Ethnicity does not account for differences in the health related quality of life of Turkish, Moroccan, and Molucan elderly in Netherlands. *Health and Quality of Life Outcomes*, 12, 138. Doi: 10.1186/s12955-014-0138-8
- Vinaccia, S., Cadena, J., Juárez, F., Contreras, F., & Anaya, J. M. (2004). Relaciones entre variables sociodemográficas, incapacidad funcional, dolor y desesperanza aprendida en pacientes con diagnóstico de artritis reumatoide. *International Journal of Clinical and Health Psychology*, 4(1), 91-103.
- Vinaccia, S., Ramírez, L., & Toro, F. (2001). Desesperanza aprendida en pacientes con artritis reumatoide: Evolución y medición de un constructo. *Revista Colombiana de Reumatología*, 8, 443-447.
- Vinaccia, S., Tobón, S., Moreno-San Pedro, E., Cadena, J., & Anaya, J. M. (2005). Evaluación de la calidad de vida en pacientes con artritis reumatoide. *International Journal of Psychology and Psychological Therapy*, 5, 45-60.
- Vinaccia, S., Fernández, F., Moreno, E., & Padilla, G. (2006). Aplicación de la versión española del cuestionario Quality of Life Measure for Rheumatoid Arthritis QOL-RA en Colombia. *Revista Colombiana de Reumatología*, 13, 284-270.
- Vinaccia, S., Quiceno, J. M., & Remor, E. (2012). Resiliencia, percepción de enfermedad, creencias y afrontamiento espiritual-religioso en relación con la calidad de vida relacionada con la salud en enfermos crónicos colombianos. *Anales de Psicología*, 28(2), 366-377.
- Vinaccia, S., Riveros, F., Quiceno, J. M., & Anaya, J. M. (en prensa). Análisis psicométrico de la versión española del cuestionario Quality of Life Measure for Rheumatoid Arthritis Scale (QOL-RA) en población colombiana. *Enfermería clínica*.
- Zhao, S., Chen, Y., & Chen, H. (2015). Sociodemographic factors associated with functional disability in outpatients with rheumatoid arthritis in Southwest China. *Clinical Rheumatology*, 34(5), 845-851. Doi: 10.1007/s10067-015-2896-z